(1) Publication number:

0 240 131

A2

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 87301522.6

O : #F....

(22) Date of filing: 23.02.87

(5) Int. Cl.³: A 61 K 31/135 A 61 K 31/205

39 Priority: 24.02.86 GB 8604528

(43) Date of publication of application: 07.10.87 Bulletin 87/41

(A) Designated Contracting States:

AT BE CH DE ES FR GB GR IT LI LU NL SE

(7) Applicant: IMPERIAL CHEMICAL INDUSTRIES PLC Imperial Chemical House Millbank London SW1P 3JF(GB)

(72) Inventor: Kraus, Hans Alfred Im Gänswasen 4 D-7050 Walblingen-Neustadt(DE)

(4) Representative: Smith, Stephen Collyer et al, Imperial Chemical Industries PLC Legal Department: Patents Po Box 6 Bessemer Road Welwyn Garden City Herts, AL7 1HD(GB)

(54) Tamoxifen for topical use.

(5) The invention concerns a new therapeutic agent for use in treating psoriasis, new compositions of the therapeutic agent for topical administration and a new method of treating psoriasis.

EP 0 240 131 A2

THE See Front Page

5

10

15

20

25

30

THERAPEUTIC AGENTS

This invention relates to a therapeutic agents for use in a new method of medical treatment and, more particularly, it relates to an agent for use in a new method for the treatment of psoriasis and to topical formulations of said agent.

Psoriasis is a common chronic inflammatory disease of the skin and in some cases, also the joints, which affects virtually all ages and which is difficult to treat successfully. The aetiology of psoriasis is largely unknown and the objective of most treatment is to suppress the abnormally rapid skin cell turnover process rather than to effect a cure. Topical treatments such as dithranol and coal tar derivatives, with or without UV irradiation, have some effect in controlling mild to moderate psoriasis, but the more severe cases require systemic treatment with cytotoxic agents such as methotrexate which have overt adverse side effects. Present treatments of psoriasis have been reviewed by R.H. Champion (British Medical Journal, 1986; 292, 1693). There is thus a need for a new therapeutic agent which is capable of controlling psoriasis without the production of unwanted side—effects.

It has now been found that administration of tamoxifen (as its citrate salt) results in the remission of psoriasis of long standing without any overt side-effects and this is the basis of the invention.

According to the present invention there is provided a method for the treatment of psoriasis affecting a warm-blooded animal which comprises the administration to said animal of an effective amount of tamoxifen or a pharmaceutically acceptable acid—addition salt thereof.

According to a further feature of the invention there is provided the use of tamoxifen, or of a pharmaceutically acceptable acid addition salt thereof, for the treatment of psoriasis affecting a warm-blooded animal such as man.

According to a yet further feature of the invention there is provided the use of tamoxifen, or of a pharmaceutically acceptable acid—addition salt thereof, for the manufacture of a novel medicament for the treatment of psoriasis affecting a warm—blooded animal such as man.

The chemical compound tamoxifen [(Z)-2-[p-(1,2-diphenylbut-1-enyl)phenoxy]ethyldimethylamine] is known to be especially useful in the treatment of hormone-dependent tumours, and especially in the treatment of breast cancer in women. Comprehensive reviews of its clinical usage are available, for example by Furr and Jordan in "Pharmacology and Therapeutics", 1984, Volume 25, pages 127-205. Tamoxifen also has utility in treating male infertility (Furr and Jordan, above, Section 8.3.1.2, page 185).

A suitable pharmaceutically acceptable acid—addition salt is, for example, the hydrochloride, hydrobromide, citrate or D-gluconate salt. The last salt is particularly suitable for the production of topical formulations of tamoxifen.

It is to be understood that "warm blooded animal" includes both male and female humans.

The tamoxifen or an acid-addition salt thereof (hereafter referred to as "the therapeutic agent") may be administered, for example orally or topically for the relief of psoriasis, and may be used in a conventional oral or topical formulation.

Topical formulations of the therapeutic agent for use in the treatment of psoriasis are novel and are provided as a further feature of the invention and comprise tamoxifen or a pharmaceutically acceptable acid—addition salt thereof (especially a generally water—soluble salt such as the D—gluconate) together with one or more dermatologically acceptable excipients.

10

15

20

25

A suitable oral formulation, is for example, a tablet or capsule, preferably a tablet containing, for example, 10, 20, 30 or 40 mg. of therapeutic agent. In general, it will be administered at a dose in the range 20 to 80 mg. per day, either as a single dose or as divided doses.

A suitable topical formulation is, for example, an ointment, cream or lotion containing up to 10% by weight of the therapeutic agent and typically in the range, for example 0.1 to 5% by weight. This will normally be applied to the skin of the patient (either to the unaffected or the psoriatic area) once or twice per day. Topical formulations may be obtained by conventional procedures and using dermatologically acceptable excipients, well known in the art.

Examples of dermatologically suitable excipients include those well known in the art of pharmacy for the production of topical formulation, such as arachis oil, non-volatile fatty alcohols, acids and/or esters (e.g. cetostearyl alcohol, cetyl alcohol, stearyl alcohol, stearic acid or cetyl esters wax); volatile alcoholic compounds [e.g. lower alkanols such as ethanol or isopropanol]; glycols and glycol ethers (e.g. propylene glycol or ethyl digol); glycerol and glycerol ethers; emulsifying agents (e.g. sorbitan stearate, sorbitan trioleate, polysorbate 60 or a mixture of polyoxyethylated oleyl or cetyl alcohol); preservatives (e.g. (1-4C)alkyl hydroxybenzoates such as methyl or propyl hydroxybenzoates); and non-volatile hydrocarbons such as soft paraffin or liquid paraffin. Examples of suitable formulations are provided in the accompanying Examples.

An agent to improve penetration through the skin, for example dimethyl sulphoxide, N-methyl-2-pyrrolidinone or l-dodecylazacycloheptan-2-one, may also conveniently be present in a topical formulation.

5

10

15

20

25

A particular ointment formulation may be prepared, for example, by dispersing the therapeutic agent as defined above in a suitable organic diluent, for example soft paraffin, optionally in the presence of an emulsifying and/or thickening agent, for example sorbitan monostearate.

A particular emulsion formulation such as a cream or lotion may be obtained for example, by adding a slurry of tamoxifen free base in ethanol or 2-propanol to an aqueous solution of D-gluconic acid to form tamoxifen gluconate in situ, which mixture is then combined with the required components of the disperse phase.

The topical formulations are conveniently provided for the treatment of psoriasis packaged in dispensers such as deformable tubes containing for example, 15 to 200 g of formulation and such filled dispensers are provided as a further feature of the invention.

One preferred formulation for oral administration contains tamoxifen citrate together with one or more pharmaceutically acceptable diluents or carrier in the form of a tablet containing 10, 20, 30 or 40 mg of active ingredient, for example, such as is commercially available under the ICI trade mark 'Nolvadex' for use in the treatment of breast cancer.

A preferred formulation for topical administration contains tamoxifen D-gluconate together with one or more dermatologically acceptable excipients in the form of an ointment, cream or lotion. In addition, the therapeutic agent may also be administered topically by spray or aerosol.

5

10

15

20

The invention is illustrated, but not limited, by the following Examples:-

EXAMPLE 1

5

10

15

25 ·

A white male patient, aged about 33 years, had been afflicted with psoriasis for about 10 years, the worst areas affected being the palms of his hands. All previous attempts to treat this condition had failed. The patient was also infertile, and treatment for the infertility was attempted by the oral administration of tamoxifen citrate (available under the trade mark 'Nolvadex' from Imperial Chemical Industries PLC) at a dose of 20 mg twice per day. After 2 months of treatment, all symptoms of the psoriasis disappeared. After a further 3 months, treatment was discontinued and the psoriasis returned 3 weeks later. Treatment with tamoxifen citrate was recommenced and the psoriasis again disappeared, and remained absent as treatment was continued. No untoward effects were observed during the period of treatment.

20 EXAMPLE 2

Clinical responses to oral treatment with tamoxifen citrate have been observed in a number of additional patients. For example a white male patient, aged about 32 years and afflicted with severe psoriasis for several years, showed a marked reduction in symptoms following administration of tamoxifen citrate orally at 20 mg twice daily. The beneficial effect was lost on withdrawing the therapeutic agent but returned with further treatment. No adverse effects were observed during treatment.

30 EXAMPLE 3 (Components are given in % w/w unless otherwise stated.)

This example describes a number of typical cream formulations for topical use and which contain tamoxifen as its D-gluconate salt.

35 For each formulation, the tamoxifen D-gluconate

salt may be obtained by mixing an ethanolic slurry of tamoxifen as the free base with an aqueous solution of D-gluconic acid, obtained by hydrolysing a 20% w/v solution of D-gluconolactone in water by heating at 70°C for 15 to 30 minutes. [Note: the minimum volume of ethanol is used and 5 ml of the aqueous D-gluconic acid solution is added per 1 g of tamoxifen free base]. Stirring is then continued until a clear solution is obtained. The required volume of this solution is then added to other excipients to produce the following cream formulations in which "active ingredient" is tamoxifen D-gluconate salt:-

FORMULATION I

5

10

	Active ingredient	0.5-2.5
15	Cetostearyl alcohol	4.0
	Cetyl esters wax	2.0
	(synthetic spermaceti)	
	Arachis oil	5.0
	* 'Brij' 72 (steareth 2)	2.5
20	** 'Brij' 78 (steareth 20)	1.5
	Methyl hydroxybenzoate	0.15
	Propyl hydroxybenzoate	0.08
	Purified water	to 100%

25 * polyethylene glycol ether of stearyl alcohol containing an average of 2 oxyethyl groups per molecule ['Brij' is a trade mark].

polyethylene glycol ether of stearyl alcohol containing an average of 20 oxyethyl groups per molecule.

PORMULATION II

	Active ingredient	0.	5-2.5
	Cetostearyl alcohol		5.0
5	Cetyl esters wax	•	3.0
	(synthetic spermaceti)	•	
	Liquid parafffin Glyceryl stearate SE		3.0
	(Self emulsifying grade)		3.0
10	'Brij' 78		1.0
	(Steareth 20)		
	Methyl hydroxybenzoate		0.15
	Propyl hydroxybenzoate		0.08
	Butylated hydroxytoluene		0.05
15	Purified water	to	100%

FORMULATION III

	Active ingredient	0.5-2.5
20	*'Arlatone' 983S	4.0
	Cetyl alcohol	2.5
	Stearic acid	1.5
25	Propylene glycol	2.0
23	Glycerin	2.0
	Methyl hydroxbenzoate	0.15
	Propyl hydroxybenzoate	0.08

Note that the property of polyethylene glycol ether derivative (containing an average of 5 oxyethyl units per molecule) of glyceryl stearate ['Arlatone' is a trade mark].

FORMULATION IV

	Active ingredient	0.5-2.5
	Cetostearyl alchol	1.0
5	Stearic acid	1.5
	*"Arlamol" E	9.0
	(PPG 15 stearyl ether)	3.0
	'Brij' 72	3.0
	(steareth 2)	
10	**'Brij' 721	2.0
	(steareth 21)	
	Propylene glycol	4.0
	Methyl hydroxybenzoate	0.15
	Propyl hydroxybenzoate	0.08
15	Purified water to 100%	

^{*} polypropylene glycol ether of stearyl alcohol containing an average of 15 2-oxypropyl groups per molecule.

20 ** polyethylene glycol ether of stearyl alcohol containing an average of 21 oxyethyl groups per molecule.

['Arlamol' is a trade mark]

25 FORMULATION V

	Active ingredient	0.5-2.5%
	Cetostearyl alcohol	5.0
•	Cetyl esters wax	2.0
30	(synthetic spermaceti)	2.0
	'Span' 60	2.0
	(sorbitan stearate)	
	'Tween' 60	2.0
	(polysorbate 60)	
35	Glycerol	1.0

Methyl hydroxybenzoate	0.15
Propyl hydroxybenzoate	0.08
Purified water to	100%

['Span' and 'Tween' are trade marks] 5

NOTES:

- (1) The methyl and propyl hydroxybenzoates are present as preservatives and may be replaced by one or more other 10 conventional preservatives.
 - (ii) The above formulations may optionally contain 0.5 to 5.0% of a penetration enhancer such as dimethyl sulphoxide.

(iii) It will be understood that the above formulations will also contain residual ethanol from the initial preparation of the active ingredient.

EXAMPLE 4 (Components are given in % w/w except where indicated)

This Example illustrates an ointment formulation suitable for topical administration:-

า	Λ
4	v

15

20	•			
	Active ingredient*		0.5	
i	Ethyl digol		5.0	
	Ethanol 96% B.P.		3.0	w/v
	Purified water		2.0	
25	'Span' 85		3.0	
	(sorbitan trioleate)			
	White soft paraffin	to	100%	

*Note: the active ingredient may either be tamoxifen D-gluconate (conveniently prepared in situ in the ethanol/ethyl digol 30 mixture) or tamoxifen citrate.

EXAMPLE 5 (Components are given in % w/w) This Example illustrates a further ointment formulation suitable for topical formulation:-

5 Active ingredient * 0.5-2.5
Light liquid paraffin 5-10
White soft paraffin to 100%

Note: the active ingredient may be, for example, finely
micronised tamoxifen citrate or finely powdered tamoxifen Dgluconate (for example such as may be obtained by freeze-drying
an aqueous solution of the salt).

CLAINS

What is claimed is -

- A pharmaceutical composition adapted for topical use which comprises as active ingredient tamoxifen or a pharmaceutically acceptable acid—addition salt thereof, together with one or more dermatologically acceptable excipients.
- 2. A composition as claimed in claim 1 which is in the form of an ointment, cream or lotion.
 - 3. A composition as claimed in claim 1 or 2 containing up to 10% by weight of the active ingredient.
 - 4. A composition as claimed in claim 1 or 2 containing 0.1 to 5% by weight of active ingredient.
- A composition as claimed in any of claims 1-4 wherein
 the active ingredient is a generally water-soluble salt of tamoxifen.
 - 6. A composition as claimed in any of claims 1-4 wherein the active ingredient is tamoxifen D-gluconate.
 - 7. A filled dispenser of topically useful pharmaceutical composition wherein the composition is as defined in any of claims 1—6 and is present in an amount of 5 to 200 g.
- 30 8. A composition as claimed in any of claims 1-6 which comprises in addition a skin penetration enhancer.

15

- 9. The use of a therapeutic agent selected from tamoxifen and the pharmaceutically acceptable acid-addition salts thereof in the manufacture of a novel medicament for use in the treatment of psoriasis.
- 10. A process for the manufacture of a novel medicament for use in the treatment of psoriasis which comprises bringing a therapeutic agent selected from tamoxifen and the pharmaceutically acceptable salts thereof into a form suitable for topical administration by combining said agent with one or more dermatologically acceptable carriers.
 - 11. A process as claimed in claim 10 wherein the therapeutic agent is selected from tamoxifen and its D-gluconate and citrate salts.

SC33777EP
SF02
SCS/MJW: 4Feb87

CLAIMS (for the contracting States Greece, Spain and Austria)

What is claimed is:-

- 1. A process for the manufacture of a novel medicament for use in the treatment of psoriasis which comprises bringing a therapeutic agent selected from tamoxifen and the pharmaceutically acceptable salts thereof into a form suitable for topical administration by combining said agent with one or more dermatologically acceptable carriers.
- 2. A process as claimed in claim 2 wherein the therapeutic agent is selected from tamoxifen and its D-gluconate and citrate salts.
 - 3. The use of a therapeutic agent selected from tamoxifen and the pharmaceutically acceptable acid—addition salts thereof in the manufacture of a novel medicament for use in the treatment of psoriasis.

SC337770EE SF02

SCS/MJW: 4Feb87

5

11 Publication number:

0 240 131 A3

12

EUROPEAN PATENT APPLICATION

(21) Application number: 87301522.6

22 Date of filing: 23.02.87

(5) Int. Cl.³: **A 61 K 31/135** A 61 K 31/205

30 Priority: 24.02.86 GB 8604528

(43) Date of publication of application: 07.10.87 Bulletin 87/41

(88) Date of deferred publication of search report: 15.11.89

Designated Contracting States:
 AT BE CH DE ES FR GB GR IT LI LU NL SE

(7) Applicant: IMPERIAL CHEMICAL INDUSTRIES PLC Imperial Chemical House Millbank London SW1P 3JF(GB)

(72) Inventor: Kraus, Hans Alfred Im Gänswasen 4 D-7050 Waiblingen-Neustadt(DE)

Representative: Smith, Stephen Collyer et al, Imperial Chemical Industries PLC Legal Department: Patents Po Box 6 Bessemer Road Welwyn Garden City Herts, AL7 1HD(GB)

(54) Tamoxifen for topical use.

(5) The invention concerns a new therapeutic agent for use in treating psoriasis, new compositions of the therapeutic agent for topical administration and a new method of treating psoriasis.



EUROPEAN SEARCH REPORT

Application Number

EP 87 30 1522

·]	DOCUMENTS CONSID	ERED TO BE RELE	VANT			
Category	Citation of document with indic of relevant passa			vant aim	CLASSIFICA APPLICATIO	TION OF THE N (Int. Cl. 4)
X	WO-A-8 503 228 (MAUV * Page 3, lines 1-19; page 5, line 22; clai	page 4, line 32	- 1-2			31/135 31/205
A	WO-A-8 401 506 (GATE * Page 2, line 23 - p page 5, line 5 *		1-8			
A	EUR. J. CANCER CLIN. no. 10, 1985, pages of the pages of	1171-1174, Pergam SE et al.: with ndrogen or ced malignant ecutive phase II	on 9-1	1		
A	CANCER TREATMENT REP 6, 1982, page 1447; al.: "Phase II trial malignant melanoma" * Whole article *	C.G. LEICHMAN et	9-1	.1		AL FIELDS D (Int. Cl.4)
A	EUROP. J. CANCER CLI no. 6, 1984, pages 8 et al.: "The effect growth of human malf vitro" * Abstract; page 807 column, line 21 - pa column, line 11; pag column, line 13 - pa column, line 39 - pag column, line 8 *	807-815; P.G. GILL of tamoxifen on dignant melanoma in 7, right-hand age 808, left-hand ge 812, right-hand age 813, left-hand	ihe 1 d	11		
	The present search report has b	een drawn up for all claims				·
	Place of search	Date of completion of	he search		Examiner	
i I	THE HAGUE	29-08-1989		GOE	TZ G.	
A:	CATEGORY OF CITED DOCUME particularly relevant if taken alone particularly relevant if combined with an document of the same category technological background: non-written disclosure intermediate document	E : earl after other D : doc L : doc	ory or principle universelve the filing date sument cited in the ment cited for our mother of the same unent cited same unent cited same unent cited same	ent, but pub e application ther reasons	lished on, or n	ing



EUROPEAN SEARCH REPORT

Application Number

EP 87 30 1522

Category	Citation of document with of relevant	n indication, where appropriate, passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
P,A	BRIT. MED. JOURNAL	., vol. 292, June .696; R.H. CHAMPION:	9-11	
	•			
1		•		
	•			
- 1				TECHNICAL FIELDS SEARCHED (Int. Cl.4)
.				
l				,
				-
	•	•		
	•			
-				
		•		
		· · · · · · · · · · · · · · · · · · ·	1	٠.
	The present search report has t	een drawn up for all claims		
	Place of search HAGUE	Date of completion of the search	000	Examiner
		29-08-1989	GOETZ	
X : partic Y : partic docum	ATEGORY OF CITED DOCUME ularly relevant if taken alone ularly relevant if combined with an nent of the same category playing hydroground	E : earlier patent d after the filing other D : document cited L : document cited	in the application for other reasons	
O : non-w	ological background written disclosure rediate document	& : member of the document	same patent family,	corresponding

THIS PAGE BLANK (USPTO)

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

BLACK BORDERS

IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

FADED TEXT OR DRAWING

BLURRED OR ILLEGIBLE TEXT OR DRAWING

SKEWED/SLANTED IMAGES

COLOR OR BLACK AND WHITE PHOTOGRAPHS

GRAY SCALE DOCUMENTS

LINES OR MARKS ON ORIGINAL DOCUMENT

REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

IMAGES ARE BEST AVAILABLE COPY.

OTHER:

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)